

Swine Influenza Assessment Criteria

Exposure Assessment Criteria

One or more of the following exposures in the 7 days prior to onset of symptoms:


- Travel to an area with current or previously documented or suspected community transmission of swine influenza (Mexico; Scotland; Israel; Canada, Spain; New Zealand; etc. — Includes transit in an airport)

OR

- Close contact with a person who is infected with swine influenza

OR

- Close contact with a person with a severe respiratory illness who has a history of travel to a foreign or domestic location with recent transmission of swine influenza



If patient meets Exposure Criteria, proceed to Clinical Assessment.

Clinical Assessment Criteria

Pre-suspect Swine Flu

- Presence of two or more of the following:
 - Temperature > 100.4° F (>38° C)
 - Chills
 - Cough
 - Sore throat
 - Nasal congestion or runny nose

Suspect Swine Flu

- Temperature > 100.4° F (>38° C)

AND

- Two or more of the following:
 - Cough
 - Chills
 - Sore throat
 - Nasal congestion or runny nose

AND

- History of travel to locations with confirmed swine flu cases

Probable Swine Flu

- Temperature > 100.4° F (>38° C)

AND

- Two or more of the following:
 - Cough
 - Chills
 - Sore throat
 - Nasal congestion/runny nose

AND

- One or more of the following:
 - History of travel to location where swine flu has been confirmed

OR

- Contact to a confirmed case of swine flu



Notify Oklahoma City-County Health Department — Epidemiology Services of any suspect or probable Swine Influenza cases prior to transport.

405-425-4437 / after hours 405-517-5536